LISTING OF THE CLAIMS

1-4. (Cancelled)

- 5. (Currently amended) An <u>isolated</u> antibody or antibody fragment thereof that activates an endogenous activity of a human erythropoietin receptor in a mammal, comprising: at least one heavy chain variable region having the amino acid sequence of SEQ ID NO:3 or antibody fragment thereof <u>comprising</u> the amino acid sequence of SEQ ID NO:3; and at least one light chain variable region having the amino acid sequence of SEQ ID NO:5 or antibody fragment thereof <u>comprising</u> the amino acid sequence of SEQ ID NO:5, wherein said antibody or antibody fragment thereof does not interact with a peptide having an amino acid sequence of SEQ ID NO:1.
- 6. (Withdrawn) An antibody or antibody fragment thereof that activates an endogenous activity of a human erythropoietin receptor in a mammal, comprising at least one heavy chain variable region having the amino acid sequence of SEQ ID NO:7 or antibody fragment thereof; wherein said antibody or antibody fragment thereof does not interact with a peptide having an amino acid sequence of SEQ ID NO:1.
- 7. (Withdrawn) An antibody or antibody fragment thereof that activates an endogenous activity of a human erythropoietin receptor in a mammal, comprising at least one light chain variable region having the amino acid sequence of SEQ ID NO:9 or antibody fragment thereof; wherein said antibody or antibody fragment thereof does not interact with a peptide having an amino acid sequence of SEQ ID NO:1.
- 8. (Withdrawn) An antibody or antibody fragment thereof that activates an endogenous activity of a human erythropoietin receptor in a mammal, said antibody comprising: at least one heavy chain variable region having the amino acid sequence of SEQ ID NO: 7 or antibody fragment thereof; and at least one light chain variable region having the amino acid sequence of SEQ ID NO: 9 or antibody fragment thereof, wherein said antibody or antibody fragment thereof does not interact with a peptide having an amino acid sequence of SEQ ID NO:1.

9-10. (Cancelled)

11. (Currently amended) An <u>isolated</u> antibody or antibody fragment thereof that activates an endogenous activity of a human erythropoietin receptor in a mammal, said antibody comprising: the amino acid sequences of at least one heavy chain variable region and at least one light chain variable region selected from the group consisting of SEQ ID NO:11/SEQ ID NO:13, SEQ ID NO:15/SEQ ID NO:17, SEQ ID

NO:19/SEQ ID NO:21, SEQ ID NO:11/SEQ ID NO:23, SEQ ID NO:11/SEQ ID NO:25, SEQ ID NO:11/SEQ ID NO:25, SEQ ID NO:33, SEQ ID NO:35/SEQ ID NO:37, SEQ ID NO:39/SEQ ID NO:41, SEQ ID NO:43/SEQ ID NO:45, SEQ ID NO:45, SEQ ID NO:47/SEQ ID NO:51/SEQ ID NO:51/SEQ ID NO:53 and SEQ ID NO:55/SEQ ID NO:57 or antibody fragment thereof comprising SEQ ID NO:51 and SEQ ID NO:53, wherein said antibody or antibody fragment thereof does not interact with a peptide having an amino acid sequence of SEQ ID NO:1.

12. (Currently amended) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one heavy chain variable region having the amino acid sequence of SEQ ID NO: 3 and at least one light chain variable region having the amino acid sequence of SEQ ID NO: 5 or antibody fragment thereof comprising SEQ ID NO:3 and SEQ ID NO:5.

13. (Cancelled)

- 14. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one heavy chain variable region having the amino acid sequence of SEQ ID NO:7 or antibody fragment thereof.
- 15. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one light chain variable region having the amino acid sequence of SEQ ID NO:9 or antibody fragment thereof.
- 16. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one heavy chain variable region having the amino acid sequence of SEQ ID NO:11 or antibody fragment thereof.
- 17. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one light chain variable region having the amino acid sequence of SEQ ID NO:13 or antibody fragment thereof.
- 18. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one heavy chain variable region having the amino acid sequence of SEQ ID NO:15 or antibody fragment thereof.

- 19. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one light chain variable region having the amino acid sequence of SEQ ID NO:17 or antibody fragment thereof.
- 20. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one heavy chain variable region having the amino acid sequence of SEQ ID NO:19 or antibody fragment thereof.
- 21. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one light chain variable region having the amino acid sequence of SEQ ID NO:21 or antibody fragment thereof.
- 22. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one light chain variable region having the amino acid sequence of SEQ ID NO:23 or antibody fragment thereof.
- 23. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one light chain variable region having the amino acid sequence of SEQ ID NO:25 or antibody fragment thereof.
- 24. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one light chain variable region having the amino acid sequence of SEQ ID NO:27 or antibody fragment thereof.
- 25. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one light chain variable region having the amino acid sequence of SEQ ID NO:29 or antibody fragment thereof.
- 26. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one heavy chain variable region having the amino acid sequence of SEQ ID NO:31 or antibody fragment thereof.
- 27. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one light chain variable region having the amino acid sequence of SEQ ID NO:33 or antibody fragment thereof.

- 28. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one heavy chain variable region having the amino acid sequence of SEQ ID NO:35 or antibody fragment thereof.
- 29. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one light chain variable region having the amino acid sequence of SEQ ID NO:37 or antibody fragment thereof.
- 30. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one heavy chain variable region having the amino acid sequence of SEQ ID NO:39 or antibody fragment thereof.
- 31. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one light chain variable region having the amino acid sequence of SEQ ID NO:41 or antibody fragment thereof.
- 32. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one heavy chain variable region having the amino acid sequence of SEQ ID NO:43 or antibody fragment thereof.
- 33. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one light chain variable region having the amino acid sequence of SEQ ID NO:45 or antibody fragment thereof.
- 34. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one heavy chain variable region having the amino acid sequence of SEQ ID NO:47 or antibody fragment thereof.
- 35. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one light chain variable region having the amino acid sequence of SEQ ID NO:49 or antibody fragment thereof.
- 36. (Currently amended) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one heavy chain variable

region having the amino acid sequence of SEQ ID NO:51 and at least one light chain variable region having the amino acid sequence of SEQ ID NO:53 or antibody fragment thereof comprising SEQ ID NO:51 and SEQ ID NO:53.

37. (Cancelled)

- 38. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one heavy chain variable region having the amino acid sequence of SEQ ID NO:55 or antibody fragment thereof.
- 39. (Withdrawn) An isolated antibody or antibody fragment thereof capable of binding to a human erythropoietin receptor in a mammal, said antibody comprising: at least one light chain variable region having the amino acid sequence of SEQ ID NO:57 or antibody fragment thereof.
- 40. (Currently amended) An isolated antibody capable of binding a human erythropoietin receptor in a mammal, said antibody comprising a heavy chain variable region comprising a continuous sequence from CDR1 through CDR3 having the amino acid sequence selected from the group consisting of: SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, and SEQ ID NO:61 and a light chain variable region comprising a continuous sequence from CDR1 through CDR3 having the amino acid sequence SEQ ID NO:62 and fragments thereof.

41. (Cancelled)

- 42. (Withdrawn) A method of activating an endogenous activity of a human erythropoietin receptor in a mammal, the method comprising the step of administering to said mammal a therapeutically effective amount of an antibody or antibody fragment thereof to activate said receptor, wherein said antibody or antibody fragment thereof does not interact with a peptide having an amino acid sequence of SEQ ID NO:1.
- 43. (Withdrawn) A method of modulating an endogenous activity of a human erythropoietin receptor in a mammal, the method comprising the step of administering to a mammal a therapeutically effective amount of the antibody or antibody fragment of claim 1 to modulate the activity of the receptor.
- 44. (Withdrawn) A method of treating a mammal suffering aplasia, the method comprising the step of administering to a mammal in need of treatment a therapeutically effective amount of an antibody or

antibody fragment thereof to activate said receptor, wherein said antibody or antibody fragment thereof does not interact with a peptide having an amino acid sequence of SEQ ID NO:1.

45. (Withdrawn) A method of treating a mammal suffering aplasia, the method comprising the step of administering to a mammal in need of treatment a therapeutically effective amount of the antibody or antibody fragment of claim 1 to modulate the activity of the receptor.

46. (Cancelled)

- 47. (Withdrawn) An isolated and purified polynucleotide sequence selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, fragments, complements, and degenerate codon equivalents thereof.
- 48. (Withdrawn) An isolated and purified amino acid sequence selected from the group consisting of: SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68 and fragments thereof.
- 49. (Currently amended) An <u>isolated</u> antibody or antibody fragment thereof that activates an endogenous activity of a human erythropoietin receptor in a mammal wherein said antibody <u>is Ab12</u>, <u>wherein Ab12 has ATCC Accession No. PTA-5554 or antibody fragment is a gamma-2 isotype</u>.

50-54. (Cancelled)

55. (Withdrawn) A method of activating an endogenous activity of a human erythropoietin receptor in a mammal, the method comprising the step of administering to a mammal a therapeutically effective amount of the antibody or antibody fragment of claim 49 to activate said receptor.

- 56. (Withdrawn) A method of modulating an endogenous activity of a human erythropoietin receptor in a mammal, the method comprising the step of administering to a mammal a therapeutically effective amount of the antibody or antibody fragment of claim 49 to modulate the activity of the receptor.
- 57. (Withdrawn) A method of treating a mammal suffering aplasia, the method comprising the step of administering to a mammal in need of treatment a therapeutically effective amount of the antibody or antibody fragment of claim 49 to activate the receptor.
- 58. (Withdrawn) A method of treating a mammal suffering aplasia, the method comprising the step of administering to a mammal in need of treatment a therapeutically effective amount of the antibody or antibody fragment of claim 49 to modulate the activity of the receptor.
- 59. (Withdrawn) A method of treating a mammal suffering anemia, the method comprising the steps of administering to a mammal in need of treatment a therapeutically effective amount of the antibody or antibody fragment of claim 49 to modulate the activity of the receptor.
- 60. (Original) A pharmaceutical composition comprising a therapeutically effective amount of the antibody or antibody fragment of claim 49 and a pharmaceutically acceptable excipient.
- 61. (New) The antibody fragment of claim 5, 11, 12, 36, 40 or 49, wherein the fragment is one selected from the group consisting of an Fab fragment, Fab' fragment, F(ab')₂ fragment and Fv fragment.